

# Reconstitution guide for BeneFIX



## PREPARATION


- Ensure product is at room temperature & wash hands thoroughly.
- Place vial on a flat surface & remove vial cap.
- Cleanse vial stopper with alcohol swab.

## RECONSTITUTION STEPS

	<p><b>STEP 1</b></p> <ul style="list-style-type: none"> <li>• Peel back cover of packaging from the plastic vial adapter and place over the vial.</li> <li>• Press down until adapter spike penetrates vial stopper.</li> <li>• Lift the package away from the adapter and discard.</li> </ul>		<p><b>STEP 2</b></p> <ul style="list-style-type: none"> <li>• Insert plunger rod into the diluent syringe and rotate to attach rod firmly.</li> </ul>
	<p><b>STEP 3</b></p> <ul style="list-style-type: none"> <li>• Break off tip cap from the diluent syringe and connect syringe to the vial adapter, turn the syringe clockwise until secured.</li> </ul>		<p><b>STEP 4</b></p> <ul style="list-style-type: none"> <li>• Inject diluent into the product vial and swirl gently to dissolve product, solution should appear clear and colourless.</li> </ul>
	<p><b>STEP 5</b></p> <ul style="list-style-type: none"> <li>• Invert the vial, draw back all the solution into the syringe.</li> <li>• Disconnect syringe by turning counter-clockwise.</li> <li>• Discard vial with adapter attached.</li> </ul>		

**Please note:**

The solution should be clear and colourless. If it is not, the solution should be discarded and a new kit should be used.



## AVAILABLE IN A RANGE OF DOSING OPTIONS:



250 IU

500 IU

1000 IU

2000 IU

## STORING BENEFIX

- BeneFIX should be kept refrigerated (at 2°C - 8°C).
- Do not freeze, to prevent damage to the diluent syringe.
- The product can be removed from the refrigerator and kept at room temperature (BELOW 30°C) for up to 6 months.
- At the end of this period, the product must not be put back in the refrigerator but must be used or discarded. It is suggested that you record the date on which the carton was removed from the refrigerator.
- Use the reconstituted solution as soon as possible after reconstitution or within 3 hours.
- The solution will be clear and colourless. Do not use BeneFIX if you notice that it is cloudy or contains visible particles.
- Do not use after the expiry date stated on the outer box and vial label.

### Haemophilia Treatment Centre Contact Details:



For further information about your condition or its treatment see your healthcare professional.



Pfizer Australia Pty Ltd, Wharf Road, West Ryde, NSW 2114. ©Registered trademark. © Pfizer Australia Pty Ltd. 2011. P7392 April 2013 PFIBEN0737. For further information please call 1800 675 229 or email [medicalaffairs.anz@pfizer.com](mailto:medicalaffairs.anz@pfizer.com).

orchard

Studio 1, 285a Crown Street,  
Surry Hills, Sydney 2010  
9339 4366

CLIENT AMEND

1

Date:	09/05/2013	Lead: Name	Last: Sarah Gleeson
Job#/Client:	Pfizer PFIBEN0737	Trim Size:	210(H) x 297(W)mm Beed: 3mm
Job Name/Ver:	PeconstitutionGuide-	Type Area:	0000(H) x 0000(W)mm

