


IDELVION® RECONSTITUTION¹


Mix2Vial™ Instructions

Step 1




Place water vial on a level surface and hold firmly. Peel the lid from the Mix2Vial blister package and pierce the WFI vial.

Step 2




Carefully remove the Mix2Vial outer package, making sure the Mix2Vial remains attached to the WFI vial.

Step 3



Hold the product vial firmly on a level surface. Pierce the powder vial (WFI flows automatically).

Incorrect Connection Methods




Step 4



Unscrew Mix2Vial and WFI vial.

Gently swirl until dissolved.

Step 5



Draw reconstituted product into syringe.

Reference: 1. IDELVION® Approved Product Information. Abbreviation: WFI, Water for Injection vial.
For detailed instructions on reconstitution and administration, see package insert or Full Product Information.

These instructions are for the Mix2Vial™ filter transfer set provided with IDELVION®



PBS Information: This product is not listed on the PBS.
National Blood Authority: This product is funded under NBA arrangements. Please refer to the NBA for details.

Before prescribing, please review Product Information available by scanning the QR code, or by contacting medicalinformation@cslbehring.com.au or 1800 642 865.

Indicated in haemophilia B patients of all ages for¹:

- Routine prophylaxis to prevent or reduce the frequency of bleeding episodes
- Control and prevention of bleeding in the perioperative setting
- Control and prevention of bleeding episodes

Contraindications: Known hypersensitivity to IDELVION®, its components, excipients, or hamster protein.¹

Precautions: Risk of allergic type hypersensitivity reactions, thrombotic complications and inhibitor development. Monitor FIX activity by one-stage clotting assay.¹ **Adverse Events:** The most frequent adverse reactions in clinical trials are headache, dizziness rash, hypersensitivity and injection site reactions. The frequency of these events is in the common category (common: $\geq 1/100$ and $< 1/10$).¹

®IDELVION is a registered trademark of CSL Limited or its affiliated companies.

Mix2Vial™ is a trademark of West Pharmaceutical Services, Inc. or a subsidiary thereof.

CSL Behring (Australia) Pty Ltd. ABN 48 160 734 761. 189–209 Camp Rd, Broadmeadows Victoria 3047.

E: customerservice@cslbehring.com.au. W: www.cslbehring.com.au

Date of preparation: June 2025. CSL Behring AU-IDL-0017



Printed on 100% recycled paper (100% post-consumer waste).



CSL Behring