## Parents' Evaluation of Developmental Status (PEDS) in Research

Title	Brief description
Waiting for speech-language pathology services: A randomised controlled trial comparing therapy, advice and device	PEDS used within a large survey. McLeod S, Davis E, Rohr K, McGill N, Miller K, Roberts A, Thornton S, Ahio N, Ivory N. Waiting for speech-language pathology services: A randomised controlled trial comparing therapy, advice and device. Int J Speech Lang Pathol. 2020 Jun;22(3):372-386. doi: 10.1080/17549507.2020.1731600. Epub 2020 May 4. PMID: 32366124. <u>Here</u>
Watch me grow integrated (WMG-I): protocol for a cluster randomised controlled trial of a web-based surveillance approach for developmental screening in primary care settings	Eapen V, Liaw S, Lingam R, et al Watch me grow integrated (WMG-I): protocol for a cluster randomised controlled trial of a web-based surveillance approach for developmental screening in primary care settings BMJ Open 2022;12:e065823. doi: 10.1136/bmjopen-2022-065823 <u>Here</u>
Sleep in the Early Years Study (2016-2017)	QUT Centre for Children's Health Research. Survey PEDS will be used to look at developmental risk in children 0-3 years. The study sample will be approx 400 children attending childcare services in Brisbane. PEDS will be administered twice on each child.
Early Childhood in Foster and Kinship Care (2015)	PEDS will be used with foster and kinship carers of young children who have not yet commenced their first year of school across Australia. The study aims to investigate participation in early learning and care programs, support for early learning offered in kinship care and foster care homes and the developmental status and wellbeing of foster and kinship care children. This research is being undertaken by the University of Melbourne and the Berry Street Childhood Institute.
New Zealand Health Survey - Child Developmental Health module (2012-2013, 2013-2014, 2014-2015, 2015-2016)	Population based survey. The New Zealand Health Survey is a cross-sectional survey of New Zealand adults and children. Parents will complete PEDS as part of a face to face interview about their children.
Early Childhood Development Assessment and Screening - Everyone's Business (2014 - 2015)	Universal surveillance of CALD families in SE Sydney. Ten supported playgroup coordinators linked to NGOs and working with CALD families will undertake developmental surveillance using PEDS.
A systematic review of the prevalence of parental concerns measured by the Parents' Evaluation of Developmental Status (PEDS) indicating developmental risk (2014)	This systematic review sought to identify the prevalence of parental concerns about their children's development, as measured by the PEDS, along with identifying associated risk factors. 37 studies were ultimately included in the review, with over 210,000 subjects in total. Nearly 14% of parents raised concerns associated with high developmental risk and overall more than 31% raised concerns associated with either high or moderate risk of developmental problems. A variety of individual, family and service level risk factors were identified. Here
A longitudinal study of late talkers and typically developing children (2012 – 2014)	Longitudinal project of toddlers with and without early language delay. The aim of this research project is to identify factors that may put children at risk of having continuing speech and language difficulties – and from an earlier age than is currently possible. The PEDS is part of the tools used to assess children at 24-30 mos of age and again at 42-48 mos. Here
Evaluation of the Cradle to Kinder and Aboriginal Cradle to Kinder programs (2013 – 2016)	This project evaluates two new ante and postnatal support services designed to provide intensive family and early parenting support to vulnerable young mothers from pregnancy through to when their child reaches four years of age. <u>here</u>



Implementing Preventive Healthcare for Young Children in General Practice (2014 - )	Part 1(2014-2018) : A pilot study educating and encouraging up to 20 General Practitioners to use P Check or in routine consultations. Part 2 (2019 - ) Extend work from Part 1 into another single prac
ACT Kindergarten Health Check - program data collection & analysis (2014 – 2018)	PEDS is being used in a State funded clinical service program, as part of a parent answered survey health check at school, at the start of the child's first year of school. PEDS data will be used for pro
Sound Start: Innovative technology to promote speech and pre-literacy skills in at risk pre-schoolers Speaking My Language: International speech acquisition in Australia (2013-15)	PEDS used to screen for suitable participants for the Sound Start intervention and as a parent repo other assessments of children's speech and language. The research is undertaken by Charles Sturt
Validation of the Child's Challenging Behaviour Scale (2011 -)	PEDS was selected as a way to stratify children with yet unidentified developmental problems. This PEDS data is not yet analysed. The research is being undertaken by a Senior Lecturer at Monash Ur
Exploring Stakeholders' Views Regarding Newborn and Infant Screening for Genetic Conditions Causing Delays in Development (2011 -)	This study explores how parents concerned about their children's development view newborn and conditions. Research is being undertaken by the Murdoch Children's Research Institute.
School-age Children Conceived via IVF and by Embryo Biopsies for Pre-implantation: Genetic Diagnosis (2011-12)	This study utilises a questionnaire that includes PEDS to explore the health and development of sc conceived via IVF and Embryo Biopsies. This research was undertaken by the Murdoch Children's R
Assessment of Medium-term Health and Development Outcomes for a Cohort of NICU survivors in Fiji (2010 -)	This study assesses use of the PEDS as a screening tool for detection of developmental delay in a g survivors (aged 2-4 years) and control newborns. The research is undertaken by The Royal Children
Children and Parents at Hospital: Improving Health and Wellbeing (CAPOH) (2010)	This study compares hospitalised children receiving usual care to those receiving an extended mod under 12 years old and experiencing stress in addition to their illness. The research was undertaken Hospital, Melbourne.
Study of Environment on Aboriginal Resilience and Child Health (SEARCH) (2008-10)	This study investigated the health and wellbeing of Aboriginal infants, children, adolescents and fa communities in New South Wales. The research is a collaboration between the Coalition for Resea Health (CRIAH) Steering Group (comprising the Aboriginal Health and Medical Research Council an Department of Health, the University of Sydney and six Aboriginal Medical Services across New Sou



PEDS as part of the Healthy Kids tice as part of a feasibility study.

completed prior to a physical ogram evaluation.

ort measure to compare with t University.

is research is still in progress and niversity.

l infant screening for genetic

chool-age children who were Research Institute.

group of newborn intensive care n's Hospital, Melbourne.

del of care. All children were en by The Royal Children's

amilies living in urban arch to Improve Aboriginal nd the Sax Institute), the NSW uth Wales. <u>Here</u>

This research followed two cohorts of children who are broadly representative of the Australian pop add to the understanding of early childhood development, inform social policy debate and help to id intervention and prevention strategies in policy areas concerning children. The research was undert Institute of Family Studies. <u>Here</u>
This study aimed to assess the feasibility of implementing baseline neurobehavioural screening follo diagnosis. The research was undertaken by the Department of Paediatrics, University of Melbourne.
This study follow Indigenous children annually with the aim of understanding what contributes to perform Indigenous children and their families. The study is undertaken by the Australian Department of Family Services and Indigenous Affairs (FaHCSIA) <u>Here</u>
This study asked teachers and parents the PEDS questions in order to identify children with speech i complements and extends findings from Wave 1 of the Longitudinal Study of Australian Children (Kir was undertaken by researchers at Charles Sturt University. <u>Here</u> <u>Here</u>
This study evaluates the assessment of the PEDS tool with two children. Research undertaken by a for Childhood Studies student, Melbourne.
This study measured the utility of PEDS and other developmental screeners with preschool-age child refugee status. Research undertaken by a Master of Public Health student, University of Western Aus
This study is a longitudinal birth cohort study based in south-western Sydney. PEDS has been used a (3½, 4, 4½ and 5 years). The study is being undertaken by researchers at the University of New South Health Care and Equity. Here
_

www.peds.org.au



opulation. The research hoped to o identify opportunities for early ertaken by the Australian

llowing childhood cancer e. <u>Here</u>

positive outcomes for amilies, Housing, Community

h impairment. The study Kindergarten Cohort). The study

a fourth year Bachelor of Early

ildren born overseas who have ustralia.

d at four time points in the study uth Wales, Centre for Primary