GRADE / YEAR

Insulin Injections

Use in conjunction with Diabetes Management Plan. This plan should be reviewed every year.

TYPE 2 INSULIN INJECTIONS SS VIC Diabetes Victoria, RCI

STUDENT NAME

DATE OF BIRTH

NAME OF SCHOOL

PHOTO

PARENT / CARER NAME

CONTACT NO.

DIABETES TREATING TEAM

HOSPITAL UR NO.

CONTACT NO.

DATE PLAN CREATED

LOW Hypoglycaemia (Hypo)

Blood Glucose Level (BGL) less than 4.0 mmol/L

SIGNS AND SYMPTOMS Pale, headache, shaky, sweaty, dizzv, drowsv, chanaes in behaviour

Note: Check BGL if hypo suspected. Symptoms may not always be obvious

DO NOT LEAVE STUDENT ALONE • DO NOT DELAY TREATMENT TREATMENT TO OCCUR WHERE STUDENT IS AT TIME OF HYPO HYPO SUPPLIES LOCATED

MILD*

Student conscious

(Able to eat hypo food)

* MILD IS COMMON

Step 1: Give fast acting carbohydrate

Step 2: Recheck BGL in 15 mins

- If BGL less than 4.0, repeat **Step 1**
- If BGL greater than or equal to 4.0, go to **Step 3**

Step 3: Give slow acting carbohydrate

If insulin is due & BGL greater than or equal to 4.0, give usual insulin dose & then eat meal immediately.

Step 3a:

Step 4: Resume usual activity when BGL 4.0 or higher

SEVERE

Student drowsy/ unconscious
(Risk of choking /

First Aid DRSABCD Stay with student

CALL AN AMBULANCE DIAL 000

Contact parent/carer when safe to do so HIGH Hyperglycaemia (Hyper)

Blood Glucose Level (BGL) greater than or equal to 15.0 mmol/L is well above target and requires additional action

SIGNS AND SYMPTOMS Increased thirst, extra toilet visits, poor concentration, irritability, tiredness

Note: Symptoms may not always be obvious

Student well

- Encourage
 1-2 glasses water per hour
- Return to usual activity
- Extra toilet visits may be required
- Re-check BGL in 2 hours

In 2 hours, if BGL still greater than or equal to 15.0,

CALL PARENT/CARER FOR ADVICE

Student unwell (e.g. vomiting)

- Contact parent/carer to collect student ASAP
- Check ketones (if strips supplied)

KETONES

If unable to contact parent/carer **and** blood ketones greater than or equal to 1.0 mmol/L

CALL AN AMBULANCE DIAL 000







Use in conjunction with Diabetes Action Plan. This plan should be reviewed every year. **TICK BOXES THAT APPLY**

INSULIN ADMINISTRATION				
INSULIN is given multiple times per day.				
The student requires an injection of insuling At home prior to school Before breakfast at before school care Lunchtime Other)			
Insulin injection minutes before				
Other diabetes medication required a				
SEE MEDICATION AUTHORITY FORM OR RELEVANT DOCUMENT				
Carbohydrate food must always be eat Location in the school where the injection			, msami mjecnom	
Is supervision required? Yes		No	Remind only	
Responsible staff will need training if they	are re	auired to:		
Administer injection (Dose as per addit		•	n provided)	
Assist Observe			,	
RESPONSIBLE STAFF Staff who have voluntarily agreed to under with diabetes care to the student.	ertake	training and p	rovide support	
with diapetes cale to the studetti.		CHICOSE	INSULIN	
STAFF MEMBER		GLUCOSE CHECKING	ADMINISTRATION / SUPERVISION	

(continues page 3)

Page 2 of 8

TYPE 2 INSULIN INJECTIONS SS VIC Diabetes Victoria, RCH, MCH 2024 V1.1

NAME__ HOSPITAL UR NO._ DATE PLAN CREATED_







INSULIN ADMINISTRATION

GLUCOSE LEVEL CHECKING

SCHOOL SETTING

A Medical Authority Form is required if school staff are to administer / supervise insulin injection / other diabetes medication.

Medication Authority Form

Ves

No

■ BEFORE / AFTER SCHOOL CARE

Before / after school care may be provided by the school, or an outside organisation. Parent / carer to obtain and complete the relevant documentation from this setting, authorising staff to administer / supervise insulin injection/other diabetes medication to their child.

GLUCOSE LEVEL CHECKING

Target range for glucose levels pre-meals: 4.0 – 7.0 mmol/L.
7.1 – 14.9 mmol/L are outside target range requiring no action.

- Glucose levels outside this target range are common.
- A glucose check should occur where the student is at the time it is required.
- Before doing a **blood glucose check** the student should wash and dry their hands.

Is the student able to do their c	own glucose level chec	:k?	
Yes	No (Support is requir	ed)	
The responsible staff member n	eeds to		
Do the check	Assist	Observe	Remind
BLOOD GLUCOSE LEVEL (BGL) TO	O BE CHECKED (tick all	those that apply)	
Anytime hypo suspected	Before snack	Before lunch	
Before activity	Before exams/tests	When feeling unwered	əll
Beginning of after-school co	are session		
Other times - please specify	,		

diabetes 💥



TYPE 2 INSULIN INJECTIONS SS VIC Diabetes Victoria, RCH, MCH 2024 V1.1



Page 3 of 8

CONTINUOUS GLUCOSE MONITORING (CGM)

- Continuous glucose monitoring consists of a small sensor that sits under the skin and measures glucose levels in the fluid surrounding the cells.
- A CGM reading can differ from a blood glucose level (BGL) reading during times of rapidly changing glucose levels e.g., eating, after insulin administration, during exercise.
- A CGM reading less than _____ mmol/L must be confirmed by a BGL check. **FOLLOW ACTION PLAN**
- Hypo treatment is based on a BGL check.
- A CGM reading above _____ mmol/L must be confirmed by a BGL check. **FOLLOW ACTION PLAN**
- If the sensor/transmitter falls out, staff to do BGL checks.

A student wearing CGM must do a blood glucose level (BGL) check: (tick all those that apply)

- Anytime hypo suspected
 When feeling unwell
- Other times please specify

USE AT SCHOOL

- Parents/carers are the primary contact for any questions regarding CGM.
- Staff are not expected to do more than the current routine diabetes care as per the student's Diabetes Action and Management plans.
- Staff do not need to put CGM apps on their personal computers, smart phones or carry receivers.
- CGM devices can be monitored remotely by family members. They should only contact the school if there is an emergency.
- The CGM sensor can remain on the student during water activities.

LOW BLOOD GLUCOSE LEVELS (Hypoglycaemia / Hypo) FOLLOW ACTION PLAN

- If the student requires more than 2 consecutive fast acting carbohydrate treatments, as per their Diabetes Action Plan, call their parent/carer. Continue hypo treatment if needed while awaiting further advice.
- All hypo treatment should be provided by parent/carer. (continues page 5)

Page 4 of 8

TYPE 2 INSULIN INJECTIONS SS VIC Diabetes Victoria, RCH, MCH 2024 V1.1

HOSPITAL UR NO. ___ DATE PLAN CREATED ___







LOW BLOOD

HIGH BLOOD

ETONES

EATING AND DRINKING

SEVERE HYPOGLYCAEMIA (HYPO) MANAGEMENT FOLLOW ACTION PLAN

Is NOT common.

DO NOT attempt to give anything by mouth to the student or rub anything onto the gums as this may lead to choking.

If the school is located more than **30 minutes** from a reliable ambulance service, then staff should discuss Glucagon injection training with the student's Diabetes Treating Team.

HIGH BLOOD GLUCOSE LEVELS (Hyperglycaemia / Hyper)

MORE THAN 15 mmol/L FOLLOW THE ACTION PLAN

KETONES FOLLOW THE ACTION PLAN

- Ketones occur most commonly in response to high glucose level and student unwell.
- Ketones are produced when the body breaks down fat for energy.
- Ketones can be dangerous.

If the student is UNWELL check ketone level if strips supplied.

FATING AND DRINKING

- Some younger students will require supervision to ensure some food is eaten.
- No food sharing.
- Seek parent/carer advice regarding foods for school parties/celebrations.
- Always allow access to water.

TYPE 2 INSULIN INJECTIONS SS VIC Diabetes Victoria, RCH, MCH 2024 V1.1

Page 5 of 8







NAME	
HOSPITAL UR NO.	
DATE PLAN CREATED_	

PHYSICAL ACTIVITY

Hypo treatment and a glucose monitoring device should always be with the student.

- Physical activity may cause glucose levels to go high or low.
- Some students may require a glucose check before, during and after physical activity.
- Some students MAY require a slow acting carbohydrate before planned physical activity.
 ADDITIONAL INFORMATION:
- Physical activity should not be undertaken if BGL less than 4.0 mmol/L.
 REFER TO THE DIABETES ACTION PLAN FOR HYPO TREATMENT
- Physical activity **should not** be undertaken if the student is **unwell**.

EXCURSIONS / INCURSIONS

It is important to plan for extracurricular activities.

- Ensure blood glucose monitor, blood glucose strips, ketone strips (if supplied), insulin device and needle and hypo food are readily available.
- Plan for meal and snack breaks.
- Always have hypo treatment available.
- Know location of toilets.

SCHOOL CAMPS

- Parents/carers need to be informed of any school camp at least 2 months prior to ensure the student's diabetes treating team can provide a Camp Diabetes Management plan and any training needs required.
- Parents/carers will need a copy of the camp menu and activity schedule.
- At least 2 responsible staff attending the camp require training to be able to support the student on camp.
- If the camp location is more than 30 minutes from a reliable ambulance service,
 Glucagon injection training is recommended.

Page 6 of 8

TYPE 2 INSULIN INJECTIONS SS VIC Diabetes Victoria, RCH, MCH 2024 V1.1









EXAMS

- Glucose level should be checked before an exam.
- Glucose level should be greater than 4.0 mmol/L before exam is started.
- Blood glucose monitor and blood glucose strips, CGM devices or smart phones, hypo treatments, and water should be available in the exam setting.
- Extra time will be required if a hypo occurs, for toilet privileges or student unwell.

APPLICATIONS FOR SPECIAL CONSIDERATION

National Assessment Program Literacy and Numeracy (NAPLAN)

Applies to Grade 3, Grade 5, Year 7, Year 9. Check National Assessment Program website - Adjustment for student with disability for further information.

Victorian Certificate of Education (VCE)

Should be lodged at the beginning of Year 11 and 12. Check Victorian Curriculum and Assessment Authority (VCAA) requirements.

EQUIPMENT CHECKLIST

Supplied by the parent/carer

- Insulin pens and pen needles Stored according to the school's Medication Policy
- Finger prick device
- Blood glucose monitor
- Blood glucose strips
- Blood ketone strips (if supplied)
- Hypo treatment
- Sharps' container
- Charging cables for diabetes management devices

DISPOSAL OF MEDICAL WASTE

- Dispose of any used pen needles in sharps container provided.
- Dispose of blood glucose and ketone strips as per the school's medical waste policy.





TYPE 2 INSULIN INJECTIONS SS VIC Diabetes Victoria, RCH, MCH 2024 V1.1



NAME	
HOSPITAL UR NO	
DATE PLAN CREATED	

Page 7 of 8

AGREEMENTS

AGREEMENTS

PARENT/CARER

Organise a meeting with school representatives to discuss implementation and sign off on your child's action and management plan.

I have read, understood, and agree with this plan.

I give consent to the school to communicate with the Diabetes Treating Team about my child's diabetes management at school. NAME FIRST NAME (PLEASE PRINT) FAMILY NAME (PLEASE PRINT) SIGNATURE DATE **SCHOOL REPRESENTATIVE** I have read, understood, and agree with this plan. NAME FIRST NAME (PLEASE PRINT) FAMILY NAME (PLEASE PRINT) ROLE Principal Vice Principal **SIGNATURE** DATE **DIABETES TREATING MEDICAL TEAM** NAME

HOSPITAL NAME

SIGNATURE

FIRST NAME (PLEASE PRINT)

Page 8 of 8

TYPE 2 INSULIN INJECTIONS SS VIC Diabetes Victoria, RCH, MCH 2024 V1.1

NAME _______HOSPITAL UR NO. ________
DATE PLAN CREATED _______



FAMILY NAME (PLEASE PRINT)

DATE



