



Victorian Children’s
Tool for Observation
and Response

5–11
years

UR NUMBER

SURNAME

GIVEN NAME(S)

DATE OF BIRTH

AFFIX PATIENT LABEL HERE ↑

Hospital _____

Frequency of Observations

Observations should be performed routinely at least 4 hourly, unless advised below							
Refer to local procedure for who can alter frequency							
Date	(e.g.) 6/4/25						
Frequency	2/24						
Name/Designation	Smith RN						

Events/Comments

Record event details, including comments, interventions and parental concerns

	Date	Time		Initial	Designation
A					
B					
C					
D					
E					
F					
G					
H					

O₂ Device NP = nasal prongs, HM = Hudson mask, HNP = humidified nasal prongs, HFNP = high flow nasal prongs

Assessment of Respiratory Distress

	Mild	Moderate	Severe
Airway	• Stridor on exertion/crying	• Some stridor at rest	• Stridor at rest
Behaviour and feeding	• Normal • Talks in sentences	• Some/intermittent irritability • Difficultly talking/crying • Difficultly feeding or eating	• Increased irritability and/or lethargy • Looks exhausted • Unable to talk or cry • Unable to feed or eat
Respiratory rate	• Mildly increased	• Respiratory rate in orange zone	• Respiratory rate in purple zone • Increased or markedly reduced respiratory rate as the child tires
Accessory muscle use	• Mild intercostal and suprasternal recession	• Moderate intercostal and suprasternal recession • Nasal flaring	• Marked intercostal, suprasternal and sternal recession
Oxygen	• No oxygen requirement	• Mild hypoxemia corrected by oxygen • Increasing oxygen requirement	• Hypoxemia may not be corrected by oxygen
Other		• May have brief apnoeas	• Gasping, grunting • Extreme pallor, cyanosis • Increasingly frequent or prolonged apnoeas

Note, not all respiratory assessment features are relevant to all conditions

Refer to your local procedure for instructions on **how** to call for assistance and escalate care

MANDATORY EMERGENCY CALL
Choose MET or other Code response

Response criteria

- Apnoea or cyanosis
- Cardiac or respiratory arrest
- Airway threat
- Prolonged convulsion
- Sudden decrease in conscious state
- Any observation in the purple zone
- 3 or more simultaneous orange zone criteria
- Staff member is very worried about the child's clinical state
- A family member is very worried about the child's clinical state

Actions required

1. Place emergency call
2. Initiate appropriate clinical care until the arrival of the emergency response team
3. Emergency response team to attend immediately, stabilise patient and/or provide advice
4. Emergency response team to document management plan

CLINICAL REVIEW RECOMMENDED

Response criteria

- Any observation in the orange zone
- Staff member is worried about the child's clinical state
- A family member is worried about the child's clinical state

Actions required

1. Initiate appropriate clinical care
 2. Consider what is usual for the child and if the trend in observations suggests deterioration
 3. Consult with nurse in charge, decide if a medical review is required
 - 4. Medical review**
 - Increase frequency of observations as indicated by the child's condition
 - If not attended within 30 minutes, escalate to emergency call
 - Medical officer to document management plan
- OR**
- 4. No medical review**
 - Document rationale & plan of care in Events/Comments

General Instructions

You MUST record baseline observations, including blood pressure, on admission and thereafter:

- At a frequency appropriate for the child's clinical state
- Whenever staff or family members are worried about the child's clinical state
- If the child is deteriorating

Level of Consciousness should be documented using the AVPU scale, **except** for children receiving sedation, where a Level of Sedation score should be recorded.

Select a Pain Assessment tool appropriate for the age, developmental level and clinical state of the child. Refer to the website and/or the RCH clinical practice guidelines for pain tools.

Show the Trend: Plot the Dot – Join the Line

This chart is specifically designed to enhance the identification of trends in vital signs. It is important to look for worsening trends and report these.

When graphing observations, place a dot in the box and connect it to the previous dot with a straight line. For blood pressure use the symbols indicated on the chart. For SpO₂ write the number in the appropriate section.

Whenever an observation falls within an orange zone or purple zone, you MUST initiate the actions required for that colour, unless a modification has been made.

Modifications — refer to your local procedure for altering calling criteria.

Level of Sedation (UMSS – University of Michigan Scoring System)

ONLY complete if sedation administered

- 0 = Awake and alert
- 1 = Minimally sedated: may appear tired/sleepy, responds to verbal conversation and/or sound
- 2 = Moderately sedated: somnolent/sleeping, easily roused with tactile stimulation or simple verbal command
- 3 = Deep sedation: deep sleep, rousable only with deep or physical stimulation
- 4 = Unrousable



The ViCTOR project is supported by the Victorian Government

Victorian Children's Tool for Observation and Response

5-11 years

Actual age:	
Weight:	

Surname:

UR:

Given name:

AFFIX PATIENT LABEL OVER PAGE

[illegible]

O ₂ Saturation (%)		(write value)
Modifications		
Purple		
Orange		
Duration <i>(maximum: 24 hrs)</i>		
Date		
Time		
Dr		
Signature		
≥94		
90–93		
≤89		
O ₂ delivery L/min or %		
Device		
Probe change		

Respiratory Rate (breaths/min)						Write ≥58					
Modifications											
Purple						55					
						52					
						49					
						46					
Orange						43					
						40					
Duration <i>(maximum 24 hrs)</i>						37					
						34					
Date						31					
						28					
Time						25					
						22					
Dr						19					
						16					
						13					
Signature						10					

[illegible]

Heart Rate (beats/min)			Write ≥170	Write ≥170
Modifications				
Purple	(e.g.) 155			165 160 155 150 145 140 135 130 125 120 115 110 105 100 95 90 85 80 75 70 65 60 55
Orange	140			
Duration <i>(maximum 24 hrs)</i>	4/24			
Date	6/4/25			
Time	1600			
Dr	Smith			
Signature	Smith			
			Write ≤50	Write ≤50

Blood Pressure $\searrow \swarrow$ (mmHg) systolic BP is the trigger			
Modifications			
Purple			
Orange			
Duration <i>(maximum 24 hrs)</i>			
Date			
Time			
Dr			
Signature			

Write ≥ 160	Write ≥ 160
155	155
150	150
145	145
140	140
135	135
130	130
125	125
120	120
115	115
110	110
105	105
100	100
95	95
90	90
85	85
80	80
75	75
70	70
65	65
60	60
55	55
50	50
Write ≤ 45	Write ≤ 45

Temperature (C°)		
Reportable limits—if applicable, refer to local procedures		
Temp ≥	(e.g.) 39.5	
Temp ≤	—	
Date	6/4/25	
Time	1800	
Dr	Smith	
Signature	<i>Smith</i>	

Write ≥40	Write ≥40
39.5	39.5
39	39
38.5	38.5
38	38
37.5	37.5
37	37
36.5	36.5
36	36
35.5	35.5
35	35

Level of Consciousness		Alert	Verbal	Pain	Unresponsive
(wake patient before scoring)	Alert				
	Verbal				
	Pain				
	Unresponsive				

[illegible][illegible]

Additional Observations (e.g. BSL, weight, capillary refill time)

[illegible]

Observations to be plotted with a dot and joined with a line (except SpO_2 and BP)