

Emergency drugs etc for child of age: 12years Expected Wt around: 40 Kg

| | | |
|--------------------------------|--------|------|
| Minimum acceptable systolic BP | 90 | mmhg |
| Heart Rate range | 60-100 | bpm |
| Respiratory Rate range | 16 | bpm |

| | | |
|--------------------------|---------|--|
| Adrenaline dose | 4.0 | mls. |
| Fluid bolus | 800 | mls of Normal Saline |
| Glucose / Dextrose (25%) | 40 | mls |
| DC Shock | 100/150 | Joules Use lower dose 1 st shock, then higher dose) |

| | | |
|----------------|---------|-------------------|
| ET tube size | 7.0 | internal diameter |
| ET tube length | 18 / 22 | cm to lip / nose |

| | | |
|------------------|-----------|--|
| Ketamine | 40 | mg (Intubation Dose – 1mg/kg) May be repeated. |
| Atropine | 0.6 | mg (Intubation Dose – 0.02mg/kg) |
| Diazepam | 8 | mg (0.2mg/kg) Beware respiratory depression |
| Thiopentone | 100 – 200 | mg (2.5-5.0mg/kg) Beware cardiovascular depression. Titrate dose |
| Suxamethonium | 50 | mg (dose varies with age) |
| Pancuronium | 4 | mg (0.1mg/kg) |
| Vecuronum | 4 | mg (0.1mg/kg) |
| Fentanyl | 80 | mcg (2mcg/kg) Beware respiratory depression. Titrate dose |
| Morphine | 4 | mg (0.1mg/kg) Beware respiratory depression. Titrate dose |
| Midazolam | 4 | mg (0.1mg/kg) Beware respiratory depression. |
| Benzylpenicillin | 2000 | mg (50mg/kg) |
| Cefotaxime | 2000 | mg (50mg/kg) |
| Mannitol | 10 | g (0.25g /kg) = 80mls of 12.5%, or 50mls of 20% |

To make an infusion of Dopamine or Dobutamine where 1ml/hr = 5mcg/kg/min add 600mg of the drug to 50mls of Normal Saline.